



Al-Oanzi ZH, Fountana S, Moonira T, Tudhope SJ,
Petrie JL, Alshawhi A, Patman G, Arden C, Reeves HL, Agius L.

[Opposite effects of a glucokinase activator and metformin on glucose-regulated gene expression in hepatocytes.](#)

Diabetes, Obesity and Metabolism (2017)

DOI: <http://dx.doi.org/10.1111/dom.12910>

Copyright:

This is the peer reviewed version of the following article, which has been published in final form at <http://dx.doi.org/10.1111/dom.12910>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Date deposited:

13/02/2017

Embargo release date:

16 February 2018

**Opposite effects of a glucokinase activator and metformin
on glucose-regulated gene expression in hepatocytes**

**Ziad H. Al-Oanzi^{1,3}, Sophia Fountana¹, Tabassum Moonira¹, Susan J Tudhope¹, John L. Petrie¹,
Ahmed Alshawi¹, Gillian Patman², Catherine Arden¹, Helen L. Reeves², Lorraine Agius^{1#}**

¹Institute of Cellular Medicine and Ageing and Health, Newcastle University,
Newcastle upon Tyne, NE2 4HH, UK

²Northern Institute for Cancer Research, Newcastle University, Newcastle-upon-Tyne, NE2 4HH, UK.

³Aljouf University, Medical Laboratory Science, P.O. B 2014, Sakaka, Saud Arabia

Correspondence to
Professor Lorraine Agius
Institute of Cellular Medicine,
The Medical School,
Newcastle upon Tyne NE2 4HH
UK
Tel +44-1912087033
E-mail: Lorraine.agius@ncl.ac.uk

Short running title: *Converse mechanisms of metformin and a GKA on liver gene regulation*

Word count: Abstract 249; Body 2799, Legends 536
5 Figures; No Tables

LA is guarantor of this work and takes responsibility for the integrity of the data and accuracy of the data analysis.

Abbreviations: AMPK, AMP-activated protein kinase; ChIP, chromatin immunoprecipitation; compound C, 6-[4-(2-Piperidin-1-ylethoxy)-phenyl]-3-pyridin-4-ylpyrazolo[1,5-a] pyrimidine (AMPK inhibitor); DG, 2-deoxyglucose; F1P, fructose 1-phosphate; G6P, glucose 6-phosphate; GKA, glucokinase activators; GKR, glucokinase regulatory protein; H6P, hexose 6-phosphate; mGPD, mitochondrial glycerol 3-phosphate dehydrogenase; T2D, Type 2 diabetes.

Abstract

Aim: Small molecule activators of glucokinase (GKAs) have been extensively explored as potential anti-hyperglycaemic drugs for type 2 diabetes (T2D). Several GKAs were remarkably effective at lowering blood glucose during early therapy but then lost their glycaemic efficacy chronically during clinical trials. We used rat hepatocytes to test the hypothesis that GKAs raise hepatocyte glucose 6-phosphate, (G6P, the glucokinase product) and down-stream metabolites with consequent repression of the liver glucokinase gene (*Gck*). We compared a GKA with metformin, the most widely prescribed drug for T2D.

Results: Treatment of hepatocytes with 25mM glucose raised cell G6P concomitantly with *Gck* repression and induction of *G6pc* (glucose 6-phosphatase) and *Pklr* (pyruvate kinase). A GKA mimicked high glucose by raising G6P and fructose-2,6-bisphosphate, a regulatory metabolite, causing a left-shift in glucose responsiveness on gene regulation. Fructose like the GKA repressed *Gck* but modestly induced *G6pc*. 2-Deoxyglucose, which is phosphorylated by glucokinase, but not further metabolised caused *Gck* repression but not *G6pc* induction, implicating the glucokinase product in *Gck* repression. Metformin counteracted the effect of high glucose on the elevated G6P and fructose 2,6-bisphosphate and on *Gck* repression, recruitment of Mlx-ChREBP to the *G6pc* and *Pklr* promoters and induction of these genes.

Conclusions: 1. Elevation in hepatocyte G6P and downstream metabolites with consequent liver *Gck* repression is a potential contributing mechanism to the loss of GKA efficacy during chronic therapy. 2. Cell metformin loads within the therapeutic range attenuate the effect of high glucose on G6P and on glucose-regulated gene expression.

Introduction

Type 2 diabetes (T2D) is a metabolic disease that manifests as a progressive rise in blood glucose which if not adequately treated causes damage to various tissues. Metformin, the most widely prescribed drug for T2D was exploited before target-directed drug discovery and its mode of action on hepatic glucose metabolism remains debated [1-3]. Its mechanism involves inhibition of mitochondrial complex I and other effects that are either dependent on activation of AMPK or independent of this kinase [2-5].

In recent years major effort has been focused on candidate drugs targeting glucokinase which is the first enzyme involved in glucose metabolism in the liver and insulin secretory cells [6]. In the liver glucokinase is regulated by adaptive binding to an inhibitor protein, GKR, which sequesters glucokinase in an inactive state in the nucleus at basal blood glucose [7]. GKR enables adaptive translocation of glucokinase to the cytoplasm after a meal in response to fructose or raised blood glucose. Small molecule glucokinase activators (GKAs) have been identified that bind to an allosteric site causing enzyme activation in hepatocytes, improved glucose tolerance and lowering of blood glucose in animal models and in human diabetes [6-9]. However despite very good initial efficacy on blood glucose control in T2D, this efficacy declined rather rapidly during clinical trials [10-12]. Whether this loss of efficacy is explained by a rise in blood triglycerides is unsettled [10-13].

High glucose concentration represses the glucokinase gene (*Gck*) in hepatocytes concomitantly with induction of glucose 6-phosphatase (*G6pc*) which catalyses the final reaction in hepatic glucose production [14,15]. This effect of glucose is often described as glucotoxicity because induction in *G6pc* and thereby hepatic glucose production by hyperglycaemia is counter-intuitive. However, repression of *Gck* and induction of *G6pc* by high glucose can be rationalized as a mechanism for preserving intracellular metabolite homeostasis at the expense of worsening hyperglycaemia [16]. In this study we tested the hypothesis that treatment of hepatocytes with a GKA represses the *Gck* gene by a mechanism linked to raised cell metabolites. We also tested whether metformin, which has opposite effects from GKAs on glucokinase translocation [17-19] also has converse effects from the GKA on cell metabolites and gene regulation at high glucose.

Materials and Methods

Materials: The GKA Ro28-1675 [20] was from Axon MedChem BV, Groningen; STK017597, an inhibitor of mGPD [21] was from Vitas-M Laboratory; S4048 a G6P transport inhibitor [22] was a gift from Sanofi Aventis.

Hepatocyte isolation and culture. Hepatocytes were isolated from male Wistar rats (body wt 200-300g) from Envigo, Bicester, UK [17]. Procedures conformed to Home Office Regulations and were approved by the University Ethics Committee. The hepatocytes were suspended in Minimum Essential Medium with 5% (v/v) calf serum and seeded on gelatin-coated plates [17]. After cell attachment the medium was replaced by serum-free medium containing 5mM glucose, 10nM dexamethasone, 1nM insulin and experiments were started after ~20h culture. Unless otherwise stated the medium contained 5mM glucose.

Hepatocyte incubations. Parallel incubations were performed for RNA extraction and metabolite determination. Glucose phosphorylation was determined with [2-³H]glucose (1μCi/ml) [17]. For determination of metformin uptake, hepatocytes were incubated with ¹⁴C-metformin (0.5μCi/ml) and incubations terminated by 2 rapid washes with 300mM sucrose. Incubations for metabolite determination were terminated 1h or 2h after substrate addition. Time courses showed maximum G6P elevation at 1-2 h with 25mM glucose.

Mouse study. Eight week old male C3H/He mice were fed Harlan Teklad (TD) diets (Harlan Labs, Madison, WI) - either an American lifestyle diet (ALIOS TD.110201: 45% calories from fat, high trans-fat with glucose/fructose drinking water) or a control diet (TD.110196: 15% calories from fat, low trans-fat without sugar) for 40 weeks *ad libitum*. Mice were culled at 48 weeks of age.

Metabolite determination. The medium was aspirated and the plates were snap-frozen in liquid nitrogen and stored at -80°C [14,15]. For hexose 6-P and adenine nucleotide determination, cells were extracted in 2.5% sulphosalicylic acid for deproteinization (10,000g, 10min) and metabolites assayed on the neutralized deproteinized extracts. Glucose 6-P (G6P) and 2-deoxyglucose 6-P (DG6P) were determined fluorometrically [15], and adenine nucleotides by a luciferase-coupled luminometric method [14]. For fructose 2,6-P₂ cells were extracted in 0.15M NaOH [15]. Except for fructose 2,6-P₂ (pmol/mg) other metabolites are expressed as nmol/mg protein.

Immunostaining and immunoactivity. Hepatocytes were fixed and immunostained for glucokinase (Santa Cruz, sc7908) [17]. For each condition, 10 fields (≥30 cells/field) were imaged for determination of the nuclear / cytoplasmic (N/C) ratio. Results are from 3 hepatocyte preparations. Immunoactivity to glucokinase and actin was as in [17] and to phospho-acetyl-CoA carboxylase (ser79-P) with an antibody from New England Biolabs (#3661).

mRNA analysis. RNA was extracted using TRIzol (Invitrogen) and cDNA was synthesized from 1μg of RNA with MMLV (Promega). Real-time RT-PCR was performed in a total volume of 10μl containing 50ng of reverse transcribed RNA and 5ng of forward and reverse primers, using a Sybr-Green

protocol (Promega) in a Roche Capillary Light Cycler [14]. The primers were as in [14,23]. Relative mRNA levels (vs control) were calculated by the delta cycle threshold method.

Chromatin immunoprecipitation (ChIP). Hepatocytes were cultured in 150cm² dishes and incubations terminated with formaldehyde (1%v/v, 10min) followed by glycine (0.125M, 5min) [23]. Binding of Mlx, ChREBP or IgG to *G6pc* and *Pklr* promoters was as in [23]. After pre-clearing, supernatants were incubated overnight (4°C) with 6µg IgG against ChREBP (Novus Biologicals, NB400-135), Mlx or control (sc-14705; sc-2027). DNA was extracted with phenol/chloroform, amplified by real-time PCR and is expressed as % input.

Statistical analysis. Data are means ± S.E.M. for the number of hepatocyte preparations indicated. Statistical analysis was by Student's *t*-test (paired or unpaired) or 2-way ANOVA with Bonferroni correction for ChIP. Correlations were by linear regression.

Results

A GKA causes a left-shift in glucose-regulated gene expression. In hepatocytes challenged with 25mM glucose several genes are either induced or repressed [24] by a mechanism linked to the raised cell metabolite levels rather than metabolic flux [14,15]. We determined the effects of a GKA [20] on glucose phosphorylation flux and hepatocyte G6P. The GKA increased phosphorylation flux and cell G6P at all glucose concentrations tested without changing ATP (Fig 1A-C). At 15mM glucose with GKA, cell G6P but not flux was higher ($P<0.05$) than with 25mM glucose alone (Fig-1B). We therefore used 15mM glucose (Fig1D-H) to compare the GKA with 2mM fructose (2F) or 25mM glucose (25G), which like the GKA promote glucokinase translocation [7,17]. High glucose (25mM) repressed *Gck* (Fig-1D) and induced *Pklr* and *G6pc* (Fig1E-F) and *Ppp1r3c* and *Txnip* (Fig1-G,H) which are targets of ChREBP-Mlx (*Pklr*,*G6pc*) or MondoA-Mlx (*Ppp1r3c*, *Txnip*) [15,23]. The GKA enhanced the effect of 15mM glucose on *Gck* repression and on induction of *Pklr* and *G6pc* and fructose (2mM) had similar but smaller effects (Fig1D-H). The GKA also raised fructose 2,6-P₂ (Fig. 1I). When the GKA was tested on mRNA expression at 5, 15 and 25mM glucose it had a greater fractional effect at 5-15mM than at 25mM glucose, indicating a left-shift in glucose-response (Fig1J-L) and *Gck* mRNA correlated negatively ($r=-0.84$, $P<0.04$) while *Pklr* and *G6pc* correlated positively ($r=0.96$, $P<0.002$) with G6P (Fig1M-O).

Repression of *Gck* by fructose correlates with G6P. Chronic feeding of animal models with fructose or sucrose has been found to either increase or decrease glucokinase protein or activity [25-27]. In mice fed a high-energy diet with fructose we found lower liver glucokinase protein (Fig-2A). Converse effects of dietary fructose in different studies [25-27] may be due to opposite effects of

hyperinsulinaemia [26] compared with fructose on liver *Gck* expression. Alternatively, the directional effect of fructose on *Gck* expression may be dose-dependent because low fructose (<2mM) promotes glucokinase translocation similar to the GKA, but high fructose (>2mM) depletes cell ATP by depleting inorganic phosphate [28] (Fig-2B). We tested the effect of 2-10mM fructose at basal or high glucose. Compared with 25 mM glucose, fructose (10mM) caused similar elevation of G6P (Fig-2C), lower elevation in fructose 2,6-P₂ (Fig-2D), similar repression of *Gck* (Fig-2E) and lower induction of *G6pc* (Fig-2F). *Gck* correlated negatively with G6P ($r=0.9$, $P<0.003$, Fig-2G) and *G6pc* correlated positively ($r=0.88$, $P<0.004$, Fig-2H) with fructose 2,6-P₂ but not with G6P ($r=0.36$, $P=0.39$). Five other ChREBP target genes responded to fructose similarly to *G6pc* (results not shown).

Repression of *Gck* by 2-deoxyglucose (DG) implicates hexose 6-P. To distinguish between a role for G6P or hexose 6-P (H6P) as distinct from fructose 2,6-P₂ in gene regulation we used 2-deoxyglucose (DG), which is phosphorylated by glucokinase to DG6P but not further metabolised by glycolysis (Fig-3A). We used S4048 to further raise H6P by inhibiting transport and hydrolysis of DG6P [15,29]. DG (5-10mM) raised H6P levels (Fig-3B) and inhibited glucose metabolism (Fig-3C) and lowered fructose 2,6-P₂ (Fig-3D). Unlike high fructose, DG did not lower ATP (Fig-3E) or increase uric acid production (not shown), which is a marker of transient ATP depletion [28]. DG repressed *Gck* (Fig. 3F) but did not induce *G6pc* or other ChREBP targets like *G6pd* and *Me1* (Fig-3G,H) and *Gck* mRNA correlated negatively ($r=0.89$, $P<0.003$) with H6P (Fig-3I).

Metformin antagonizes gene regulation by high glucose. Previous studies reported that GKAs stimulate [17] and high metformin concentrations (> 1mM) inhibit [18,19] glucokinase translocation. We confirmed that the GKA (Ro 28-1675) promotes translocation of glucokinase and that metformin (≥ 1 mM) inhibits translocation at high glucose (Fig-S1). We next tested whether these metformin concentrations have opposite effects from the GKA on gene regulation at high glucose in either the presence of insulin (Fig-4A-D) or its absence (Fig-4E,F). In the presence of insulin, metformin induced *Gck* at 5mM and 25mM glucose (Fig-4A) and had no effect on *Pklr*, *G6pc*, and *Txnip* at 5mM glucose but counteracted the induction by 25mM glucose (Fig-4B-D). In the absence of insulin metformin likewise counteracted induction of *G6pc* and *Txnip* by high glucose but had no effect at 5mM glucose (Fig-4E,F). This indicates counter-effects of metformin on gene induction by high glucose that are independent of insulin signalling. We tested the effects of compound C (CC), an inhibitor of AMPK and various other protein kinases [30], which blocks the metformin induction of FGF21 [31]. This inhibitor partly attenuated ($\leq 25\%$) but did not abolish the metformin effect at high glucose (Fig-5G,H), implicating a different mechanism from FGF21 induction [31]. Fructose 2,6-P₂

7which is involved in ChREBP-translocation by high glucose [15,23] was lowered by metformin at 5mM and 25mM glucose (Fig-5I) and recruitment of Mlx and ChREBP to the *Pklr* and *G6pc* promoters by high glucose was also inhibited by metformin (Fig-5J,K).

Low metformin lowers G6P at high glucose. When assessing whether the exposure to metformin in cellular studies corresponds to therapeutic doses in man [33] both incubation time and cellular metformin content are relevant. Metformin is cationic at physiological pH and is transported across the cell membrane by organic cation transporters [2] and distributes in cells (“accumulates”) [34] in accordance with transmembrane potential [2,32]. The time lag in metformin efficacy in cellular studies is in part due to slow equilibration [32]. In man blood metformin levels after an oral dose peak at 3 h (at 8 to 23 μ M for a 0.5g to 1.5 g metformin dose) and then decline with a mean elimination half-life of 5 h and a urinary excretion $t_{1/2}$ of 20 h [33]. In mice given an equivalent oral metformin load (50mg/kg) blood metformin levels peak at 0.5 to 1h (\sim 28 μ M) and decline more rapidly than in man [33,34]. The peak hepatic content of metformin in mice is higher than the coincident blood peak in the periphery and portal vein, as expected [2,32] and was 286 to 137 nmol / g liver, which approximates to 1 to 2 nmol/mg protein (based on 10^8 hepatocytes per g and 1.7 mg protein per million hepatocytes) [34]. We used 14 C-metformin to determine the cellular metformin load as a function of incubation time (Fig. 5A,B). At 2 to 4 hours the hepatocyte metformin load was 1.3 and 2.3 nmol / mg cell protein with 0.1mM and 0.2mM metformin respectively, but \geq 10 nmol/ mg with 1mM metformin (Fig. 5A,B). We infer that with 0.1 to 0.2 mM metformin the hepatocyte metformin load at 2 h approximates the peak hepatic level in mice after a therapeutic dose. Based on the time course we used a 2h pre-incubation with metformin (0.2-1mM) at basal glucose followed by 1h incubation at 5mM or 25mM glucose to determine the effects of metformin on G6P and adenine nucleotides. Metformin (0.2-1mM) attenuated the elevation in G6P at 25mM glucose (Fig-5C) and had negligible effect on total adenine nucleotides below 1mM (Fig-5D-F) although changes in free concentrations cannot be excluded [32]. Metformin (\geq 0.1mM) also lowered G6P at high glucose in the presence of S4048 (Fig-5G,H) which raises G6P levels \sim 10-fold by inhibiting G6P transport and hydrolysis [15,29]. Two mechanisms implicated at low metformin loads are activation of AMPK [4] and inhibition of mGPD [5], which is involved in transfer of NADH equivalents from the cytoplasm to mitochondria. The metformin lowering of G6P (Fig-5G,H) was not mimicked by either STK017597 an inhibitor of mGPD [21] or by A769662 an AMPK activator [35] that mimics the phosphorylation of acetyl-CoA carboxylase by 0.2-0.5mM metformin (Fig-5I). Metformin (0.2-1mM) caused concentration-dependent induction of *Gck* at 5mM and 25mM glucose and attenuated the induction of *G6pc* and *Pklr* by high glucose but did not affect these genes at basal glucose (Fig. 5J-L).

This indicates that metformin loads $\geq 1\text{-}2$ nmol/mg have converse effects on gene regulation from the GKA (Fig.1).

Discussion

The low liver glucokinase activity and *Gck* mRNA in poorly controlled T2D [36,37] is generally attributed to impaired *Gck* induction by insulin [37]. However animal studies using either hyperglycaemic clamps [38] or chronic treatment of hyperglycaemia with inhibitors of renal glucose reabsorption [39,40] support a mechanism for repression of glucokinase by hyperglycaemia [7]. Key questions are whether this repression is due to glucose *per se* [40] or to raised intracellular metabolites derived from glucose [7]. The latter would implicate potential links between antihyperglycaemic drugs that affect glucose metabolites and liver *Gck* expression. We show in this study that a GKA mimics the repression of *Gck* by high glucose whereas metformin has the opposite effect.

We identified three conditions mimicking *Gck* repression by 25mM glucose in hepatocytes: (i) the GKA at basal or moderate glucose (5-15mM); (ii) fructose, 2-10mM; (iii) 2-deoxyglucose. In all conditions *Gck* mRNA correlated inversely with hexose 6-P levels (Figs 1-3). The 2-deoxyglucose experiments implicate the raised hexose 6-P in the *Gck* repression, but do not exclude involvement of other metabolites. We showed previously that in hepatocytes incubated with high glucose, selective lowering of fructose 2,6-P₂, (without lowering G6P) by overexpression of a kinase-deficient bisphosphatase variant of PFKFB2 raised *Gck* expression [15]. This implicates a potential dual role for hexose 6-P (present study) and fructose 2,6-P₂ [15] in *Gck* repression. The rapid (within 4 hours) “metabolite repression” of liver *Gck* in hepatocytes concurs with acute repression of liver *Gck* mRNA in rats treated with S4048 which raises hepatic G6P but not blood glucose [29] or with a liver-selective GKA before a glucose tolerance test [41]. A key question is how one reconciles this mechanism with the chronic effects of GKAs over a time scale of months [10-12] or of a high-fructose diet [25-27]. Repression of glucokinase protein or activity by fructose-containing diets in some but not other studies [25-27] could be explained by the opposing effects of *Gck* repression by fructose (Fig 2) versus induction by secondary hyperinsulinaemia [26]. The half-life of glucokinase protein is 4.6 days and thus two orders of magnitude longer than for the mRNA [14]. If metabolite repression of liver *GCK* were the only adaptive mechanism to the GKA, then loss of liver glucokinase protein would be expected within 4-7 days as occurs during fasting [7]. However, loss of glycaemic efficacy of GKAs that target both liver and pancreatic islets occurred after 4 weeks in man [10,11], although in some preclinical studies this manifested earlier [41,42]. For GKAs that target both liver

and pancreatic islets, the latter effect leads to raised insulin release into the portal vein which induces the liver *GCK* gene [7]. This could explain why the glycaemic response to GKAs improves progressively for the first 4 weeks before it declines [10,11]. The first phase may represent the insulin induction of liver *GCK* and the second metabolite repression. The possibility that tachyphylaxis [10,11] is also due to loss of GKA efficacy in pancreatic islets cannot be excluded. However liver selective GKAs are effective in lowering blood glucose [43] and more so initially than chronically [41]. Furthermore in a mouse model of an activating *Gck* mutation, liver glucokinase protein is markedly decreased despite preserved glucose-induced insulin secretion and in the absence of raised lipids [44]. This suggests that down-regulation of liver glucokinase protein can occur in the absence of steatosis with a *Gck* activating mutation. Absence of steatosis was also noted in patients with *GCK* activating mutations [45]. Cumulatively the present data suggests that metabolite repression of the liver *GCK* gene is a plausible component of tachyphylaxis to GKAs [10,11]. However in the clinical setting for GKAs that also target the pancreas, tachyphylaxis in islets might be an additional mechanism.

Metformin was exploited as an anti-hyperglycaemic agent for T2D long before its mechanism of action was explored. Despite clear evidence for AMPK activation by metformin [4], also at low concentrations [46] there remain cogent arguments for AMPK-independent mechanisms [3,5]. Here we focused on the metformin effects on gene regulation at high glucose because this is the defining feature of T2D and because previous work with high metformin ($\geq 2\text{mM}$) reported inhibition of glucokinase translocation [18,19] which is an opposite response to GKAs [17]. We report here that low metformin concentrations ($\geq 0.1\text{mM}$) in incubation times of 2-4 h that lead to cell metformin loads of 1-2 nmol/mg attenuate G6P levels in conditions of elevated glucose and have opposite effects on glucose-responsive genes from the GKA. Lowering of liver G6P in rats at 4h after treatment with metformin has been observed previously [32]. In the present study the lowering of G6P by metformin was not blocked by an inhibitor of mGPD [21] and it was not mimicked by A769662 an AMPK activator [35]. However, in the absence of studies on AMPK-deficient cells we cannot firmly exclude a role for AMPK activation.

A key finding from this study is that metformin lowers G6P in hepatocytes in conditions of elevated glucose and at a cellular load of metformin that occurs during the absorptive phase in mouse liver after an oral therapeutic dose of metformin [34]. One limitation of our study is that after a 2 hour incubation of hepatocytes with low extracellular metformin concentrations, the intracellular concentrations are quasi static and therefore not the same as the dynamic changes that occur in liver after an oral therapeutic dose of metformin [34] when blood metformin levels rise rapidly and decline with a half-life of 2 to 6 hours [33,34]. Given the differences in dynamic changes between

cellular models (Fig. 5) and the liver in vivo [34] it is uncertain whether therapeutic doses of metformin will attenuate hepatic G6P in man as occurs in animal models [32] and whether such changes if they occur in man would be confined to the absorptive phase of metformin when blood levels are highest [33,47].

Acknowledgements:

Funded by Diabetes UK (Ref 11/0004231; 13/0004701) and by British Council Fellowship support (BAE PDSR) to ZAO.

Author contributions: ZHA designed experiments and researched and analysed the data. SF, TM, SJH, JLP, AA, GP performed experiments and analysed the data. CA and HLR provided samples and contributed to the discussion. LA directed the study, had access to the data, wrote the paper and is the guarantor of the work. We thank: Dr. Dieter Schmolli, Sanofi Aventis for the gift of S4048.

Conflicts of interest None.

References

1. Tahrani AA, Bailey CJ, Del Prato S, Barnett AH. Management of type 2 diabetes: new and future developments in treatment. *Lancet*. 2011;378:182-97
2. Baur JA, Birnbaum MJ. Control of gluconeogenesis by metformin: does redox trump energy charge? *Cell Metab*. 2014;20:197-199
3. Foretz M, Guigas B, Bertrand L, Pollak M, Viollet B. Metformin: from mechanisms of action to therapies. *Cell Metab*. 2014;20:953-966
4. He L, Wondisford FE. Metformin action: concentrations matter. *Cell Metab*. 2015;21:159-162
5. Madiraju AK, Erion DM, Rahimi Y, Zhang XM, Braddock DT, Albright RA, Prigaro BJ, Wood JL, Bhanot S, MacDonald MJ, Jurczak MJ, Camporez JP, Lee HY, Cline GW, Samuel VT, Kibbey RG, Shulman GI. Metformin suppresses gluconeogenesis by inhibiting mitochondrial glycerophosphate dehydrogenase. *Nature*. 2014;510:542-6.
6. Matschinsky FM. GKAs for diabetes therapy: why no clinically useful drug after two decades of trying? *Trends Pharmacol Sci*. 2013;34:90-99
7. Agius L. Hormonal and Metabolite Regulation of Hepatic Glucokinase. *Annu Rev Nutr*. 2016;36:389-415
8. Sarabu R, Berthel SJ, Kester RF, Tilley JW. Novel glucokinase activators: a patent review (2008 - 2010). *Expert Opin Ther Pat*. 2011;21:13-33
9. Filipinski KJ, Pfeifferkorn JA. A patent review of glucokinase activators and disruptors of the glucokinase--glucokinase regulatory protein interaction: 2011-2014. *Expert Opin Ther Pat*. 2014;24:875-891

10. Meininger GE, Scott R, Alba M, et al. Effects of MK-0941, a novel glucokinase activator, on glycemic control in insulin-treated patients with type 2 diabetes. *Diabetes Care*. 2011;34:2560-2566
11. Wilding JP, Leonsson-Zachrisson M, Wessman C, Johnsson E. Dose-ranging study with the glucokinase activator AZD1656 in patients with type 2 diabetes mellitus on metformin. *Diabetes Obes Metab*. 2013;15:750-759
12. Kiyosue A, Hayashi N, Komori H, Leonsson-Zachrisson M, Johnsson E. Dose-ranging study with the glucokinase activator AZD1656 as monotherapy in Japanese patients with type 2 diabetes mellitus. *Diabetes Obes Metab*. 2013;15:923-930
13. Agius L. Lessons from glucokinase activators: the problem of declining efficacy. *Expert Opin Ther Pat*. 2014;24:1155-1159
14. Arden C, Petrie JL, Tudhope SJ, et al. Elevated glucose represses liver glucokinase and induces its regulatory protein to safeguard hepatic phosphate homeostasis. *Diabetes*. 2011;60:3110-3120
15. Arden C, Tudhope SJ, Petrie JL, et al. Fructose 2,6-bisphosphate is essential for glucose-regulated gene transcription of glucose-6-phosphatase and other ChREBP target genes in hepatocytes. *Biochem J*. 2011;443:111-123
16. Agius L. Dietary carbohydrate and control of hepatic gene expression: mechanistic links from ATP and phosphate ester homeostasis to the carbohydrate-response element-binding protein. *Proc Nutr Soc*. 2016;75:10-18
17. Brocklehurst KJ, Payne VA, Davies RA, et al. Stimulation of hepatocyte glucose metabolism by novel small molecule glucokinase activators. *Diabetes*. 2004;53:535-541
18. Guigas B, Bertrand L, Taleux N, et al. 5-Aminoimidazole-4-carboxamide-1-beta-D-ribofuranoside and metformin inhibit hepatic glucose phosphorylation by an AMP-activated protein kinase-independent effect on glucokinase translocation. *Diabetes*. 2006;55:865-874
19. Mukhtar MH, Payne VA, Arden C, et al. Inhibition of glucokinase translocation by AMP-activated protein kinase is associated with phosphorylation of both GKRP and 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase. *Am J Physiol* 2008;294:R766-R774
20. Grimsby J, Sarabu R, Corbett WL, et al. Allosteric activators of glucokinase: potential role in diabetes therapy. *Science*. 2003;301:370-373
21. Orr AL, Ashok D, Sarantos MR, Ng R, Shi T, Gerencser AA, Hughes RE, Brand MD. Novel inhibitors of mitochondrial sn-glycerol 3-phosphate dehydrogenase. *PLoS One*. 2014;9:e89938.
22. Herling AW, Burger H, Schubert G, Hemmerle H, Schaefer H, Kramer W. Alterations of carbohydrate and lipid intermediary metabolism during inhibition of glucose-6-phosphatase in rats. *Eur J Pharmacol*. 1999;386:75-82.
23. Petrie JL, Al-Oanzi ZH, Arden C, et al. Glucose induces protein targeting to glycogen in hepatocytes by fructose 2,6-bisphosphate-mediated recruitment of MondoA to the promoter. *Mol Cell Biol*. 2013;33:725-738
24. Ma L, Robinson LN, Towle HC. ChREBP*MLx is the principal mediator of glucose-induced gene expression in the liver. *J Biol Chem*. 2006;281:28721-28730
25. Coate KC, Kraft G, Moore MC, Smith MS, et al. Hepatic glucose uptake and disposition during short-term high-fat vs. high-fructose feeding. *Am J Physiol* 2014;307:E151-160
26. Francini F, Castro MC, Schinella G, et al. Changes induced by a fructose-rich diet on hepatic metabolism and the antioxidant system. *Life Sci*. 2010;86:965-971
27. Sakamoto E, Seino Y, Fukami A, et al. Ingestion of a moderate high-sucrose diet results in glucose intolerance with reduced liver glucokinase activity and impaired glucagon-like peptide-1 secretion. *J Diabetes Investig*. 2012;3:432-440

28. Petrie JL, Patman GL, Sinha I, et al. The rate of production of uric acid by hepatocytes is a sensitive index of compromised cell ATP homeostasis. *Am J Physiol*. 305:E1255-E1265, 2013
29. van Dijk TH, van der Sluijs FH, Wiegman CH, et al. Acute inhibition of hepatic glucose-6-phosphatase does not affect gluconeogenesis but directs gluconeogenic flux toward glycogen in fasted rats. A pharmacological study with the chlorogenic acid derivative S4048. *J Biol Chem*. 2001;276:25727-25735
30. Bain J, Plater L, Elliott M, Shpiro N, Hastie CJ, McLauchlan H, Klevernic I, Arthur JS, Alessi DR, Cohen P. The selectivity of protein kinase inhibitors: a further update. *Biochem J*. 2007;408:297-315
31. Nygaard EB, Vienberg SG, Ørskov C, Hansen HS, Andersen B. Metformin stimulates FGF21 expression in primary hepatocytes. *Exp Diabetes Res*. 2012;2012:465282
32. Owen MR, Doran E, Halestrap AP. Evidence that metformin exerts its anti-diabetic effects through inhibition of complex 1 of the mitochondrial respiratory chain. *Biochem J*. 2000;348:607-614
33. Graham GG, Punt J, Arora M, Day RO, Doogue MP, Duong JK, Furlong TJ, Greenfield JR, Greenup LC, Kirkpatrick CM, Ray JE, Timmins P, Williams KM. Clinical pharmacokinetics of metformin. *Clin Pharmacokinet*. 2011;50:81-98
34. Wilcock C, Bailey CJ. Accumulation of metformin by tissues of the normal and diabetic mouse. *Xenobiotica*. 1994;24:49-57
35. Vincent EE, Coelho PP, Blagih J, Griss T, Viollet B, Jones RG. Differential effects of AMPK agonists on cell growth and metabolism. *Oncogene*. 2015;34:3627-39
36. Basu A, Basu R, Shah P, et al. Effects of type 2 diabetes on the ability of insulin and glucose to regulate splanchnic and muscle glucose metabolism: evidence for a defect in hepatic glucokinase activity. *Diabetes*. 2000;49:272-283
37. Haeusler RA, Camastra S, Astiarraga B, et al. Decreased expression of hepatic glucokinase in type 2 diabetes. *Mol Metab*. 2014;4:222-226
38. Massillon D, Chen W, Barzilai N, et al. Carbon flux via the pentose phosphate pathway regulates the hepatic expression of the glucose-6-phosphatase and phosphoenolpyruvate carboxykinase genes in conscious rats. *J Biol Chem*. 1998;273:228-234
39. Brichard SM, Henquin JC, Girard J. Phlorizin treatment of diabetic rats partially reverses the abnormal expression of genes involved in hepatic glucose metabolism. *Diabetologia*. 1993;36:292-298
40. Ueta K, O'Brien TP, McCoy GA, et al. Glucotoxicity targets hepatic glucokinase in Zucker diabetic fatty rats, a model of type 2 diabetes associated with obesity. *Am J Physiol Endocrinol Metab*. 2014;306:E1225-E1238
41. Erion DM, Lapworth A, Amor PA, Bai G, et al. The hepatoselective glucokinase activator PF-04991532 ameliorates hyperglycemia without causing hepatic steatosis in diabetic rats. *PLoS One*. 2014;9:e97139
42. De Ceuninck F, Kargar C, Charton Y, Goldstein S, Perron-Sierra F, Ilic C, Caliez A, Rolin JO, Sadlo M, Harley E, Vinson C, Ktorza A. S 50131 and S 51434, two novel small molecule glucokinase activators, lack chronic efficacy despite potent acute antihyperglycaemic activity in diabetic mice. *Br J Pharmacol*. 2013;169:999-1010.
43. Lloyd DJ, St Jean DJ Jr, Kurzeja RJ, Wahl RC, et al. Antidiabetic effects of glucokinase regulatory protein small-molecule disruptors. *Nature*. 2013;504:437-40.
44. Pino MF, Kim KA, Shelton KD, Lindner J, Odili S, Li C, Collins HW, Shiota M, Matschinsky FM, Magnuson MA. Glucokinase thermolability and hepatic regulatory protein binding are essential factors for predicting the blood glucose phenotype of missense mutations. *J Biol Chem*. 2007;282:13906-16.

45. Challis BG, Harris J, Sleight A, Isaac I, Orme SM, Seevaratnam N, Dhatariya K, Simpson HL, Semple RK. Familial adult onset hyperinsulinism due to an activating glucokinase mutation: implications for pharmacological glucokinase activation. *Clin Endocrinol (Oxf)*. 2014;81:855-61.
46. Meng S, Cao J, He Q, Xiong L, Chang E, Radovick S, Wondisford FE, He L. Metformin activates AMP-activated protein kinase by promoting formation of the $\alpha\beta\gamma$ heterotrimeric complex. *J Biol Chem*. 2015;290:3793-802.
47. Christensen MM, Højlund K, Hother-Nielsen O et al. Steady-state pharmacokinetics of metformin is independent of the OCT1 genotype in healthy volunteers. *Eur J Clin Pharmacol*. 2015;7:691-7.

Figure 1 Repression of *Gck* and induction of *G6pc* by a GKA

A-C: Hepatocytes were incubated for 2h in MEM containing 10nM insulin and 5, 15, or 25mM glucose (G) +/- 10 μ M GKA for determination of glucose phosphorylation (nmol/2h per mg) and cell G6P and ATP (nmol/mg protein). Means \pm SEM, n= 4-6, * $P<0.05$ effect of GKA.

D-I. Incubations for mRNA analysis (D-H) or fructose 2,6-bisphosphate (I) were for 4h without (open bars) or with (filled bars) 10nM insulin and the concentrations (mM) of glucose (G), fructose (F) indicated. Means \pm SEM, n= 3-6, # $P<0.05$ relative to 15mM glucose.

J-L. Incubations for mRNA analysis were for 4h with 10nM insulin and 5,15 or 25 mM glucose (G) +/- 10 μ M GKA. Means \pm SEM, n= 3, * $P<0.05$ effect of GKA. M-O Correlations of mRNA vs G6P.

Figure 2 Repression of *Gck* by fructose.

A. Glucokinase immunoactivity in livers of mice fed a high-energy fructose-containing diet (HEFD), Means \pm SEM, n=11, $P<0.02$ relative to control.

B. Fructose is metabolised to fructose 1-P (F1P), which activates glucokinase by dissociating it from GKRP. At high fructose (>2mM) accumulation of F1P causes depletion of inorganic phosphate (Pi) and ATP by limiting oxidative phosphorylation (OP).

C-F. Incubations were with 10nM insulin and 5 or 25mM glucose (5G/25G) and 2-10mM fructose (F) for either 2h (C,D) for G6P and fructose 2,6-P₂ or 4h (E,F) for mRNA analysis. Means \pm SEM, n= 3-8. * $P<0.05$ effect of fructose; # $P<0.05$ relative to 25G. G,H Correlation of mRNA versus metabolite.

Figure 3. Repression of *Gck* by 2-deoxyglucose

A. 2-Deoxyglucose (DG) is phosphorylated by glucokinase to DG6P which can be hydrolysed in the endoplasmic reticulum but is not metabolised by glycolysis. S4048 inhibits transport and hydrolysis of G6P and DG6P (H6P).

B-I. Incubations were with 10nM insulin and 5 or 25mM glucose (G), +/-2 μ M S4048 and 2-deoxyglucose (DG, 5 or 10mM), for either 2h (B-E) for metabolite and flux analysis or 4h (F-H) for mRNA. Means \pm SEM, n= 4-8, * $P<0.05$ relative to 5G; # $P<0.05$ relative to 25G. I, correlation of mRNA vs H6P.

Figure 4 Metformin induces *Gck* and counteracts gene induction by high glucose.

A-H. Incubations were for 4h with 10nM insulin (A-D) or without insulin (E-H) with 5mM or 25mM glucose (G) and 1mM or 5mM metformin and -/+ 8μM compound C (CC) for mRNA analysis. Means ± SEM, n=3-8, **P*<0.05 effect of metformin, #*P*<0.05 effect of compound C.

I. Fructose 2,6-P₂ (pmol/mg protein) Means ± SEM, n=3, **P*<0.05 effect of metformin.

J-K. Chromatin immunoprecipitation for binding of Mlx and ChREBP to the Pklr and G6pc promoters after 4h incubation expressed as % input. Means±SEM, n= 3, **P*<0.05 effect of metformin.

Figure 5 Metformin lowers G6P at high glucose

A,B. Cell accumulation of ¹⁴C-metformin: A, time course; B, 4h incubation. Means±SEM, n= 4.

C-I. Hepatocytes were pre-incubated for 2h with metformin and other inhibitors or activators (2μM S4048, 10-40μM SKT017597, GPI); 5-20μM A769662, A76) in MEM with 5mM glucose and then for 1h at 5 or 25 mM glucose. Means±SEM, n=4, **P*<0.05 effect of metformin. I. Immunoactivity to ACC-P (ser-79) and GAPDH, **P*<0.05 relative to 20μM A769662.

J-L. Incubations were for 4h with the additions indicated for mRNA analysis. Means±SEM, n=4, **P*<0.05 effect of metformin.

Fig 1




Figure 1 Repression of *Gck* and induction of *G6pc* by a GKA

A-C: Hepatocytes were incubated for 2h in MEM containing 10nM insulin and 5, 15, or 25mM glucose (G) -/+ 10μM GKA for determination of glucose phosphorylation (nmol/2h per mg) and cell G6P and ATP (nmol/mg protein). Means \pm SEM, n = 4-6, * $P < 0.05$ effect of GKA.

D-I. Incubations for mRNA analysis (D-H) or fructose 2,6-bisphosphate (I) were for 4h without (open bars) or with (filled bars) 10nM insulin and the concentrations (mM) of glucose (G), fructose (F) indicated. Means \pm SEM, n = 3-6, # $P < 0.05$ relative to 15mM glucose.

J-L. Incubations for mRNA analysis were for 4h with 10nM insulin and 5,15 or 25 mM glucose (G) -/+ 10 μM GKA. Means \pm SEM, n = 3, * $P < 0.05$ effect of GKA. M-O Correlations of mRNA vs G6P.

Fig 2




Figure 2 Repression of *Gck* by fructose.

A. Glucokinase immunoactivity in livers of mice fed a high-energy fructose-containing diet (HEFD), Means \pm SEM, $n=11$, $P<0.02$ relative to control.

B. Fructose is metabolised to fructose 1-P (F1P), which activates glucokinase by dissociating it from GKRP. At high fructose (>2 mM) accumulation of F1P causes depletion of inorganic phosphate (Pi) and ATP by limiting oxidative phosphorylation (OP).

C-F. Incubations were with 10nM insulin and 5 or 25mM glucose (5G/25G) and 2-10mM fructose (F) for either 2h (C,D) for G6P and fructose 2,6-P₂ or 4h (E,F) for mRNA analysis. Means \pm SEM, $n=3-8$. * $P<0.05$ effect of fructose; # $P<0.05$ relative to 25G. G,H Correlation of mRNA versus metabolite.

Fig 3




Figure 3. Repression of *Gck* by 2-deoxyglucose

A. 2-Deoxyglucose (DG) is phosphorylated by glucokinase to DG6P which can be hydrolysed in the endoplasmic reticulum but is not metabolised by glycolysis. S4048 inhibits transport and hydrolysis of G6P and DG6P (H6P).

B-I. Incubations were with 10nM insulin and 5 or 25mM glucose (G), -/+2μMS4048 and 2-deoxyglucose (DG, 5 or 10mM), for either 2h (B-E) for metabolite and flux analysis or 4h (F-H) for mRNA. Means ± SEM, n= 4-8, **P*<0.05 relative to 5G; #*P*<0.05 relative to 25G. I, correlation of mRNA vs H6P.

Fig 4




Figure 4 Metformin induces *Gck* and counteracts gene induction by high glucose.

A-H. Incubations were for 4h with 10nM insulin (A-D) or without insulin (E-H) with 5mM or 25mM glucose (G) and 1mM or 5mM metformin and -/+ 8 μ M compound C (CC) for mRNA analysis. Means \pm SEM, n=3-8, * $P < 0.05$ effect of metformin, # $P < 0.05$ effect of compound C.

I. Fructose 2,6-P₂ (pmol/mg protein) Means \pm SEM, n=3, * $P < 0.05$ effect of metformin.

J-K. Chromatin immunoprecipitation for binding of Mlx and ChREBP to the *Pklr* and *G6pc* promoters after 4h incubation expressed as % input. Means \pm SEM, n= 3, * $P < 0.05$ effect of metformin.

Fig 5




Figure 5 Metformin lowers G6P at high glucose

A,B. Cell accumulation of ^{14}C -metformin: A, time course; B, 4h incubation. Means \pm SEM, n= 4.

C-I. Hepatocytes were pre-incubated for 2h with metformin and other inhibitors or activators (2 μM S4048, 10-40 μM SKT017597, GPi); 5-20 μM A769662, A76) in MEM with 5mM glucose and then for 1h at 5 or 25 mM glucose. Means \pm SEM, n=4, * P <0.05 effect of metformin. I. Immunoactivity to ACC-P (ser-79) and GAPDH, * P <0.05 relative to 20 μM A769662.

J-L. Incubations were for 4h with the additions indicated for mRNA analysis. Means \pm SEM, n=4, * P <0.05 effect of metformin.

